

# Primary aortoenteric fistula following endovascular aortic repair due to type II endoleak

James T. McPhee, MD,<sup>a</sup> David I. Soybel, MD,<sup>b</sup> Robert K. Oram, MD,<sup>c</sup> and Michael Belkin, MD,<sup>a</sup>  
*Boston, Mass; and Dover, NH*

An 84-year-old female was lost to follow-up after endovascular aneurysm repair at another hospital with known type II endoleak. She later presented with presyncope and hematemesis. A referral center esophagogastroduodenoscopy showed possible duodenal diverticulum. She had recurrent symptoms and repeat computed tomography scan showed air within the aortic sac. At our center, she underwent stent graft explantation and axillofemoral reconstruction for a primary aortoenteric fistula. She was discharged and is doing well 5 months postoperatively. A high degree of suspicion for aortoenteric fistula is imperative in any patient with upper gastrointestinal hemorrhage after open or endovascular abdominal aortic aneurysm repair. (*J Vasc Surg* 2011;54:1164-6.)

Primary aortoenteric fistula (AEF) is a direct communication between the native aorta and the overlying bowel. It is an exceedingly rare and difficult diagnosis to make, and most are associated with abdominal aortic aneurysms. Secondary AEF complicates <2% of open abdominal aortic aneurysm (AAA) repairs.<sup>1</sup> Less is known about the true incidence and presentation of AEF formation following endovascular aortic repair (EVAR). Existing reported cases of AEF following EVAR have described both infectious and hemorrhagic manifestations with multiple etiologic factors.<sup>2-15</sup> We describe a unique presentation of a primary AEF following EVAR manifested by a subacute gastrointestinal hemorrhage.

## CASE REPORT

In the spring of 2006, an 84-year-old female presented to a referring hospital with abdominal pain and an 8-cm infrarenal, fusiform AAA. She was urgently treated for a symptomatic aneurysm by endovascular means. She underwent uneventful aortic endografting with a 28-mm AneuRx graft (Medtronic, Minneapolis, Minn) with bilateral iliac extension limbs (16 mm × 11.5 mm and 16 mm × 8.5 mm).

Interval computed tomography angiogram (CTA) 2 weeks postoperatively was notable for a type II endoleak with stable aneurysm diameter. The patient then refused further imaging and ultimately was lost to follow-up, having last been seen in good condition at her 5-month postoperative visit.

Four years later, this now 88-year-old patient presented to medical attention at an outside facility with recurrent episodes of



Fig 1. Endoscopic photograph of duodenal erosion with recent hemorrhage.

nausea, dizziness, and anemia over a 3-month period, requiring blood transfusion (hematocrit 22%) on one occasion. During this evaluation, she had a witnessed episode of hematemesis. Esophagogastroduodenoscopy (EGD) demonstrated an “unusual” posterior duodenal ulcer with no stigmata of recent bleeding, and a CT scan revealed an approximately 9.5-cm aneurysm sac. Concern was raised for AEF, and she was transferred to a tertiary care facility for further management.

At the referral center, the patient suffered no further recurrent episodes of gastrointestinal bleeding. She underwent a CT angiogram that demonstrated a type II endoleak, and repeat EGD was interpreted as consistent with a duodenal diverticulum (Fig 1). The patient was treated with aggressive acid reduction therapy and discharged to home on hospital day 5.

Three weeks later, we accepted this patient in transfer to our facility in hemodynamically normal condition for recurrent symptoms of dizziness, melena, and anemia (hematocrit 22%). The clinical history and abdominal CT findings of air within the aneurysm sac (Fig 2) confirmed the suspicion for AEF. After extensive discussion with the patient and her family in regard to the risk involved with aortic graft explantation, she was taken urgently to

From the Division of Vascular and Endovascular Surgery,<sup>a</sup> and the Division of General Surgery,<sup>b</sup> Brigham and Women's Hospital, Boston; and the Vascular Center of New Hampshire, Dover.<sup>c</sup>

Competition of interest: none.


Reprint requests: James T. McPhee, MD, Division of Vascular and Endovascular Surgery, Brigham and Women's Hospital, 75 Francis St., Boston, Mass 02115 (e-mail: [jtmcphee@partners.org](mailto:jtmcphee@partners.org)).

The editors and reviewers of this article have no relevant financial relationships to disclose per the JVS policy that requires reviewers to decline review of any manuscript for which they may have a competition of interest.

0741-5214/\$36.00

Copyright © 2011 by the Society for Vascular Surgery.

doi:10.1016/j.jvs.2011.04.049



**Fig 2.** Noncontrast computed tomography scan with air bubbles within aortic sac suggestive of aortoduodenal fistulization.

the operating room for repair in alignment with her expressed wishes.


In the operating room, she first underwent a right axillofemoral bypass graft. Abdominal exploration revealed marked periaortic inflammation with an obvious site of fistula formation between the third portion of the duodenum and the anterior aortic sac (Fig 3). Proximal infrarenal aortic and distal common iliac artery control was established. Upon opening of the aneurysm, there was no evidence of purulence within the aortic sac. While there was bile staining on the wall of the aortic sac and on the mural thrombus, there was none on the stent graft itself. A single posteromedial lumbar vessel with robust pulsatile back-bleeding into the sac was identified, confirming type II endoleak. This was controlled with a suture ligature. There was no evidence of bleeding from the proximal or distal endograft seal zones or between graft components. Likewise, the fabric of the graft was intact. The aperture to the duodenal fistula was identified from within the aneurysm sac on the anterior wall. The graft was explanted without difficulty, and the proximal and distal aortic stumps were closed in two layers. After copious irrigation with antibiotic solution, the duodenum was repaired with 3-0 silk sutures in a single-layer inverting “Gambee” stitch.<sup>16</sup> The duodenal repair sites as well as the proximal aortic stump were buttressed with a pedicled flap of vascularized omentum.

She was treated with a 7-day course of broad-spectrum antibiotics (vancomycin, ciprofloxacin, metronidazole, fluconazole). She was discharged to inpatient rehabilitation on postoperative day 21 after a prolonged ileus. At her most recent 5-month postoperative visit, she continues to do well. She ambulates with a walker, and a sterile right groin seroma is being managed expectantly.

## DISCUSSION

AEF formation following EVAR remains an extremely rare phenomenon with available literature limited only to case reports and small series.<sup>1-15</sup> Clinical suspicion remains the sine qua non of making this diagnosis due to its relative rarity and lack of a definitive diagnostic modality.

The current report is novel in that our patient presented with an intermittent subacute gastrointestinal hemorrhage related to aortoenteric fistulization that lasted nearly 3 months without rapid enough bleeding to manifest true hemodynamic compromise. We believe a persistent type II endoleak caused progressive aortic sac enlargement with



**Fig 3.** Intraoperative photograph. Depicted is cephalad reflection of the third portion of the duodenum with forceps within the large posterior duodenal defect.

ultimate erosion into the third portion of the duodenum. This likely manifested as a relatively low-pressure gastrointestinal (GI) bleed related to a back-bleeding lumbar artery with intermittent self-limited hemorrhage as opposed to the systemic aortic pressure seen in the more classic presentations of AEF. Additionally, the patient never manifested any systemic infectious signs or symptoms to indicate a primary graft infection. This case is best characterized as a primary AEF, as native aortic wall eroded into the duodenum from ongoing aneurysmal enlargement as opposed to a foreign body erosion/or infectious cause.

Prior case reports of AEF following endograft placement have represented more “classical” presentations of aortoenteric fistulization. Saratzis and colleagues recently reported five cases of aortoduodenal fistulization with varying endograft types. All of their patients presented with brisk hematemesis and hemodynamic compromise.<sup>8</sup> Earlier reports also denote the primary symptom on presentation commonly as brisk bleeding and/or shock.<sup>7,11</sup> In contrast, several other early series report infectious symptomatology.<sup>3,4</sup> A recent systematic review of reported cases of AEF following aortic endograft placement lacks specific details about each case, but did demonstrate that most patients will present with more than one complaint.<sup>17</sup>

The etiology of AEF formation varies widely in the reported cases as well. Reported cases account for AEF formation due to eroded translumbar and transluminal aortic coils<sup>10,11</sup> and type I endoleaks with sac expansion,<sup>2,8,9,18</sup> as well as graft kinking/erosion and primary fabric failure.<sup>3,6,7</sup>

This case additionally highlights the importance of patient selection for endografting as well as the concept of patient “buy-in” to maintain surveillance of these endografts postimplantation. In this case, the patient refused further radiographic imaging 6 weeks post-EVAR with a known type II endoleak and presented 4

years later with a significantly enlarged aortic sac and AEF. Persistent type II endoleaks and incomplete surveillance are both markers for poor outcomes and increased mortality following EVAR.<sup>19,20</sup>

EVAR is the predominant treatment modality for intact infrarenal AAA repair in the United States.<sup>21,22</sup> While AEF formation in the EVAR population is extremely rare, it may present in diverse ways due to myriad etiologies. As is the case after open aortic surgery, a high index of suspicion must be maintained when these patients present with gastrointestinal bleeding to avoid delay in timely intervention.

## REFERENCES

1. Busuttill SJ, Goldstone J. Diagnosis and management of aortoenteric fistulas. *Sem Vasc Surg* 2001;14:302-11.
2. del Moral LR, Alonso SF, Kiuri SS, Caballero DF, Heredero AF, Nistal MG, et al. Aortoenteric fistula arising as a complication of endovascular treatment of abdominal aortic aneurysm. *Ann Vasc Surg* 2009;23:255e13-15.
3. Hausegger KA, Tiesenhhausen K, Karaic R, Tauss J, Koch G. Aortoduodenal fistula: a late complication of intraluminal exclusion of an infrarenal aortic aneurysm. *J Vasc Med Biol* 1999;10:747-50.
4. Makar R, Reid J, Pherwani AD, Johnston LC, Hannon RJ, Lee B, et al. Aorto-enteric fistula following endovascular repair of abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 2000;20:589-90.
5. Ohki T, Veith FJ, Shaw P, Lipsitz E, Suggs WD, Wain RA, et al. Increasing incidence of mid-term and long-term complications after endovascular graft repair of abdominal aortic aneurysms: a note of caution based on a 9-year experience. *Ann Surg* 2001;234:323-35.
6. Janne d'Othee B, Soula P, Ota P, Cahill M, Joffe F, Cerene A, et al. Aortoduodenal fistula after endovascular stent-graft of an abdominal aortic aneurysm. *J Vasc Med Biol* 2000;31:190-5.
7. Norgren L, Bengt J, Engellau L. Aortoenteric fistula caused by a ruptured stent-graft: a case report. *J Endovasc Surg* 1998;5:269-72.
8. Saratzis N, Saratzis A, Melas N, Ktenidis K, Kiskinis D. Aortoduodenal fistulas after endovascular stent-graft repair of abdominal aortic aneurysms: single center experience and review of the literature. *J Endovasc Ther* 2008;15:4410-8.
9. Abou-Zamzam AM, Bianchi C, Mazraany W, Teruya TH, Hopewell J, Vannix RS, et al. Aortoenteric fistula development following endovascular abdominal aortic aneurysm repair: a case report. *Ann Vasc Surg* 2003;17:119-22.
10. Bertges DJ, Vilella ER, Makaroun MS. Aortoenteric fistula due to endoleak coil embolization after endovascular AAA repair. *J Endovasc Ther* 2003;10:130-5.
11. Elkouri S, Blair JF, Therasse E, Oliva VL, Bruneau L, Soulez G, et al. Aortoduodenal fistula occurring after type II endoleak treatment with coil embolization of the aortic sac. *J Vasc Surg* 2003;37:461-4.
12. Parry DJ, Waterworth A, Kessel D, Robertson I, Berridge DC, Scott DJ, et al. Endovascular repair of an inflammatory abdominal aortic aneurysm complicated by aortoduodenal fistulation with an unusual presentation. *J Vasc Surg* 2001;33:874-9.
13. Ruby BJ, Cogbill TH. Aortoduodenal fistula 5 years after endovascular abdominal aortic aneurysm repair with the Ancure stent graft. *J Vasc Surg* 2007;45:834-6.
14. Kar B, Dougherty K, Reul GJ, Krajcer Z. Aortic stent-graft infection due to a presumed aortoenteric fistula. *J Endovasc Ther* 2002;9:901-6.
15. French JR, Simring DV, Merrett N, Thursby P. Aorto-enteric fistula following endoluminal abdominal aortic aneurysm repair. *ANZ J Surg* 2004;74:397-99.
16. Paul S, Soybel DI, Zinner MJ. Stomach and duodenum: operative procedures. In: *Maingot's abdominal operations*. 11th ed. New York: McGraw Hill Companies, Inc; 2007. p. 387.
17. Bergqvist D, Björck M, Nyman R. Secondary aortoenteric fistula after endovascular aortic interventions: a systematic literature review. *J Vasc Interv Radiol* 2008;19:163-5.
18. Ratchford EV, Morrissey NJ. Aortoenteric fistula: a late complication of endovascular repair of an inflammatory abdominal aortic aneurysm. *Vasc Endovasc Surg* 2006;40:487-91.
19. Jones JE, Atkins MD, Brewster DC, Chung TK, Kwolek CJ, LaMuraglia GM, et al. Persistent type 2 endoleak after endovascular repair of abdominal aortic aneurysm is associated with adverse late outcomes. *J Vasc Surg* 2007;46:1-8.
20. Jones WB, Taylor SM, Kalbaugh CA, Joels CS, Blackhurst DW, Langan EM 3rd, et al. Lost to follow-up: a potential under-appreciated limitation of endovascular aneurysm repair. *J Vasc Surg* 2007;46:434-40.
21. Schwarze ML, Shen Y, Hemmerich J, Dale W. Age-related trends in utilization and outcome of open and endovascular repair of abdominal aortic aneurysm in the United States: 2001-2006. *J Vasc Surg* 2010;51:1348-52.
22. Hill JS, MCPhee JT, Messina LM, Ciocca RG, Eslami MH. Regionalization of abdominal aortic aneurysm repair: evidence of a shift to high-volume centers in the endovascular era. *J Vasc Surg* 2008;48:29-36.

Submitted Mar 5, 2011; accepted Apr 21, 2011.